

RECEIVED  
CENTRAL FAX CENTER  
SEP 28 2006

**AMENDMENTS TO THE CLAIMS**

The listing of claims will replace all prior versions, and listings of claims in the application:

**LISTING OF THE CLAIMS**

1-22. (Canceled)




23. (Previously Presented) A method for treating a genetic disorder, or condition or disease in a patient in need of treatment, comprising:








administering an effective amount of a compound comprising a cholesterol group or derivative thereof having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a straight chain polyamine; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group.

24. (Previously Presented) The method according to claim 23 wherein the cholesterol group or derivative thereof is cholesterol.


25. (Previously Presented) The method according to claim 23 wherein the cholesterol group is linked to the head group *via* a carbamoyl linkage.

26. (Previously Presented) The method according to claim 23 wherein the compound is selected from compounds of the formula















where Chol denotes a group of the formula











27. (Previously Presented) The method according to claim 23 wherein the compound is selected from compounds of the formula




28. (Previously Presented) The method according to claim 23 wherein the compound is selected from compounds of the formula






29. (Previously Presented) The method according to claim 23 wherein the compound is of the formula



30. (Previously Presented) The method according to claim 23 wherein the compound is of the formula

NCCNC(CCN)CCCCNCCCNC(=O)OC

33. (Previously Presented) The method according to claim 32, wherein the cationic lipid compound is in admixture with a nucleotide sequence.

35. (Previously Presented) The method according to claim 34, wherein the cationic liposome is in admixture with a nucleotide sequence.









administering an effective amount of a compound selected from the group consisting of cationic lipid compounds, cationic liposomes formed from cationic lipid compounds, cationic lipid compounds in admixture with a nucleotide sequence, cationic liposomes, formed from a cationic lipid compound, in admixture with a nucleotide sequence, and combinations thereof,



the compound comprising a cholesterol group or derivative thereof having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a straight chain polyamine; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group.

37. (Previously Presented) The method according to claim 36 wherein the cholesterol group or derivative thereof is cholesterol.


38. (Previously Presented) The method according to claim 36 wherein the cholesterol group is linked to the head group *via* a carbamoyl linkage.

39. (Previously Presented) The method according to claim 36 wherein the compound is selected from compounds of the formula














where Chol denotes a group of the formula









40. (Previously Presented) The method according to claim 36 wherein the compound is selected from compounds of the formula









41. (Previously Presented) The method according to claim 36 wherein the compound is selected from compounds of the formula









42. (Previously Presented) The method according to claim 36 wherein the compound is of the formula



43. (Previously Presented) The method according to claim 36 wherein the compound is of the formula



44. (Previously Presented) The method according to claim 36 wherein the compound is of the formula



45. (Previously Presented) A method for treating a genetic disorder, or condition or disease in a patient in need of treatment, comprising:

administering an effective amount of a composition, the composition comprising:

- i. a compound selected from the group consisting of cationic lipid compounds, cationic liposomes formed from a cationic lipid compound, and combinations thereof, the compound






comprising a cholesterol group or derivative thereof having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a straight chain polyamine; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group, and






- ii. a pharmaceutical, and optionally a pharmaceutically acceptable diluent, carrier or excipient.

46. (Previously Presented) The method according to claim 45 wherein the cholesterol group or derivative thereof is cholesterol.


47. (Previously Presented) The method according to claim 45 wherein the cholesterol group is linked to the head group via a carbamoyl linkage.

48. (Previously Presented) The method according to claim 45 wherein the compound is selected from compounds of the formula














where Chol denotes a group of the formula












49. (Previously Presented) The method according to claim 45 wherein the compound is selected from compounds of the formula






50. (Previously Presented) The method according to claim 45 wherein the compound is selected from compounds of the formula







51. (Previously Presented) The method according to claim 45 wherein the compound is of the formula



52. (Previously Presented) The method according to claim 45 wherein the compound is of the formula



53. (Previously Presented) The method according to claim 45 wherein the compound is of the formula



54. (Previously Presented) A method for treating a genetic disorder or condition or disease in a patient in need of treatment, comprising:




administering an effective amount of a composition comprising a compound selected from the group consisting of cationic lipid compounds, cationic lipid compounds in admixture with a nucleotide sequence, cationic liposomes (formed from a cationic lipid compound) in admixture with or associated with a nucleotide sequence, and combinations thereof;








the compound comprising a cholesterol group or derivative thereof having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a straight chain polyamine; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group.

55. (Previously Presented) The method according to claim 54 wherein the cholesterol group or derivative thereof is cholesterol.


56. (Previously Presented) The method according to claim 54 wherein the cholesterol group is linked to the head group *via* a carbamoyl linkage.

57. (Previously Presented) The method according to claim 54 wherein the compound is selected from compounds of the formula















where Chol denotes a group of the formula











58. (Previously Presented) The method according to claim 54 wherein the compound is selected from compounds of the formula




59. (Previously Presented) The method according to claim 54 wherein the compound is selected from compounds of the formula









60. (Previously Presented) The method according to claim 54 wherein the compound is of the formula



61. (Previously Presented) The method according to claim 54 wherein the compound is of the formula



62. (Previously Presented) The method according to claim 54 wherein the compound is of the formula



63. (Previously Presented) The method according to claim 54 wherein the composition further comprises a pharmaceutical.

64. (Previously Presented) The method according to claim 63, wherein the composition further comprises a pharmaceutically acceptable diluent, carrier or excipient.

65. (Previously Presented) A method for the treatment of a genetic disorder or condition or disease in a patient in need thereof, comprising administering a cationic lipid compound, the compound comprising a cholesterol group or derivative thereof having linked thereto a head group, wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a polyamine group which is a straight chain polyamine group; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group.

66. (Previously Presented) A method for the treatment of a genetic disorder or condition or disease in a patient in need thereof, comprising administering a cationic liposome formed from a cationic lipid compound, the compound comprising a cholesterol group having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a polyamine group which is a straight chain polyamine group; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group.

67. (Previously Presented) A method for the treatment of a genetic disorder or condition or disease in a patient in need thereof, comprising administering a cationic

lipid compound in admixture with a nucleotide sequence, the compound comprising a cholesterol group or derivative thereof having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a polyamine group which is a straight chain polyamine group; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group.

68. (Previously Presented) A method for the treatment of a genetic disorder or condition or disease in a patient in need thereof, comprising administering a cationic liposome in admixture with a nucleotide sequence, wherein the cationic liposome is formed from a cationic lipid compound, the compound comprising a cholesterol group having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a polyamine group which is a straight chain polyamine group; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group.

69. (Previously Presented) A method for the treatment of a genetic disorder or condition or disease in a patient in need thereof, comprising administering a pharmaceutical composition comprising

- (i) a cationic lipid compound, the compound comprising a cholesterol group or derivative thereof having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a polyamine group which is a straight chain polyamine group; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group; and
- (ii) a pharmaceutical and, optionally, a pharmaceutically acceptable diluent, carrier or excipient.

70. (Previously Presented) A method for treatment of a genetic disorder or condition or disease in a patient in need thereof, comprising administering a pharmaceutical composition comprising:

- (i) a cationic liposome formed from a cationic lipid compound, the compound comprising a cholesterol group having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a polyamine group which is a straight chain polyamine group; further wherein two or

more of the amine groups of the polyamine group are separated by an ethylene group;  
and

(ii) a pharmaceutical and, optionally, a pharmaceutically acceptable diluent, carrier or excipient.

71. (New) A method for delivering a therapeutic agent to one or more cells comprising:


administering an effective amount of a composition comprising:








(i) a compound comprising a cholesterol group or derivative thereof having linked thereto a head group; wherein the head group is more positive than the head group of DC-Chol; further wherein the head group is a straight chain polyamine; further wherein two or more of the amine groups of the polyamine group are separated by an ethylene group; and (ii) a nucleic acid.

72. (New) The method according to claim 71 wherein the cholesterol group or derivative thereof is cholesterol.


73. (New) The method according to claim 71 wherein the cholesterol group is linked to the head group *via* a carbamoyl linkage.

74. (New) The method according to claim 71 wherein the compound is selected from compounds of the formula





where Chol denotes a group of the formula



75. (New) The method according to claim 71, wherein the compound is selected from the group consisting of cationic lipid compounds, cationic liposomes formed from cationic lipid compounds, and combinations thereof.